

Godambe (1960), Geyer (2004), Neyman (1959), Pearson (1894) Rao (1948), Pearson (1900), Rao (2002), Bera and Biliias (2001).

## References

- Bera, A. K. and Y. Biliias (2001). Rao's score, Neyman's  $C(\alpha)$  and Silvey's LM tests: an essay on historical developments and some new results. *Journal of Statistical Planning and Inference* 97, 9–44.
- Geyer, C. J. (2004). Probability and statistics using nelson-style nonstandard analysis. Presentation at the symposium held to celebrate the 25th anniversary of the Department of Statistics, University of Washington, Seattle.
- Godambe, V. P. (1960). An optimum property of regular maximum likelihood estimation. *The Annals of Mathematical Statistics* 31, 1208–1211.
- Neyman, J. (1959). Optimal asymptotic tests of composite statistical hypotheses. In U. Grenander (Ed.), *Probability and Statistics. The Harald Cramér Volume*, pp. 213–234. Stockholm/New York: Almqvist & Wiksell/Wiley.
- Pearson, K. (1894). Contribution to the mathematical theory of evolution. *Philosophical Transactions of the Royal Society of London* 185, 71–110.
- Pearson, K. (1900). On the criterion that a given system of observations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. *Philosophical Magazine* 50, 157–175.
- Rao, C. R. (1948). Large sample tests of statistical hypotheses concerning several parameters with applications to problems of estimation. *Proceedings of the Cambridge Philosophical Society* 44, 50–57.
- Rao, C. R. (2002). Karl Pearson chi-square test. The dawn of statistical inference. In C. Huber-Carol, N. Balakrishnan, M. S. Nikulin, and M. Mesbah (Eds.), *Goodness-of-fit Tests and Model Validity*, Chapter 2, pp. 9–24. Boston: Birkhäuser.